

STABLE $\mathbb{A}^1$ -CONNECTEDNESS

NGUYEN LE DANG THI

ABSTRACT. We prove in this note a stabilized version of a conjecture on $\mathbb{A}^1$ -connectedness of A. Asok and B. Dorant [AD09, Conj. 2.18]. For the stabilized version of this conjecture, we introduce the notion of stable $\mathbb{A}^1$ -connectedness, which can be seen as the stabilization of $\mathbb{A}^1$ -connectedness. The notion of stable $\mathbb{A}^1$ -connectedness is in connection to the existence of zero cycles of degree one rather than rational points.

1. INTRODUCTION

Let k be a base field. We denote by $\mathcal{H}(k)$ the unstable $\mathbb{A}^1$ -homotopy category constructed in [MV01]. A k -space $\mathcal{X}$ is $\mathbb{A}^1$ -connected, if $\pi_0^{\mathbb{A}^1}(\mathcal{X}) \xrightarrow{\cong} \pi_0^{\mathbb{A}^1}(\mathrm{Spec} k) = \mathrm{Spec} k$, where we denote by $\pi_0^{\mathbb{A}^1}(\mathcal{X})$ the Nisnevich sheaf

$$\pi_0^{\mathbb{A}^1}(\mathcal{X}) = a_{\mathrm{Nis}}[U \mapsto \mathrm{Hom}_{\mathcal{H}(k)}(U, \mathcal{X})].$$

It is conjectured by A. Asok and B. Dorant that:

Conjecture 1.1. [AD09, Conj. 2.18] *Let X be a smooth $\mathbb{A}^1$ -connected scheme and $U \subset X$ an open subscheme such that $\mathrm{codim}(X \setminus U) \geq 2$. If k is an infinite field, then U is also $\mathbb{A}^1$ -connected.*

Remark this conjecture is answered by A. Asok in [A12, Prop. 3.8]. In this note, we establish the stabilized version of this conjecture. Denote by $\mathcal{SH}(k)$ resp. $\mathcal{SH}_s(k)$ the $\mathbb{P}^1$ -stable resp. the S^1 -stable homotopy category.

Definition 1.2. *Let $\mathcal{X} \in \mathrm{Spc}(k)$ be a k -space. $\mathcal{X}$ is stable $\mathbb{A}^1$ -connected, if the canonical map of sheaves*

$$\pi_0^{st}(\Sigma_{\mathbb{P}^1}^{\infty} \mathcal{X}_+) \rightarrow \pi_0^{st}(\Sigma_{\mathbb{P}^1}^{\infty} \mathrm{Spec} k_+)$$

is an isomorphism.

In [A12], A. Asok characterized the notion of unstable $\mathbb{A}^1$ -connectedness (cf. [A12, Thm. 4.15]). A smooth proper k -scheme X is $\mathbb{A}^1$ -connected if and only if the canonical map of sheaves $\mathbf{H}_0^{\mathbb{A}^1}(X) \rightarrow \mathbf{H}_0^{\mathbb{A}^1}(\mathrm{Spec} k) = \mathbb{Z}$ is an isomorphism. Remark that if X is $\mathbb{A}^1$ -connected, then $X(k) \neq \emptyset$. One has Hurewicz functors on homotopy categories $\mathcal{SH}_s(k) \rightarrow D_{\mathbb{A}^1}^{eff}(k)$ resp. $\mathcal{SH}(k) \rightarrow D_{\mathbb{A}^1}(k)$, which induce morphisms of sheaves $\pi_i^{st}(\Sigma_s^{\infty} \mathcal{X}_+) \rightarrow \mathbf{H}_i^{\mathbb{A}^1}(\mathcal{X})$ resp. $\pi_i^{st}(\Sigma_{\mathbb{P}^1}^{\infty} \mathcal{X}_+) \rightarrow \mathbf{H}_i^{st\mathbb{A}^1}(\mathcal{X})$ for a k -space $\mathcal{X}$. The stable Hurewicz theorem said that (cf. [AH11, Thm. 2.3.8]) the Hurewicz morphism

$$\pi_0^{st}(\Sigma_{\mathbb{P}^1}^{\infty} \mathcal{X}_+) \rightarrow \mathbf{H}_0^{st\mathbb{A}^1}(\mathcal{X})$$

Date: 27. 08. 2012.

1991 Mathematics Subject Classification. 14F22, 14F42.

Key words and phrases. stable $\mathbb{A}^1$ -homotopy.

is an isomorphism contravariantly functorial in $\mathcal{X}$. In particular, if $\mathcal{X} = \operatorname{Spec} k$, one has $\mathbf{K}_0^{MW} \cong \mathbf{H}_0^{st\mathbb{A}^1}(\mathbf{1}_k)$. So the characterization for the notion of stable $\mathbb{A}^1$ -connectedness is: A k -space $\mathcal{X}$ is stable $\mathbb{A}^1$ -connected if and only if $\mathbf{H}_0^{st\mathbb{A}^1}(\mathcal{X}) \xrightarrow{\cong} \mathbf{K}_0^{MW} = \mathbf{GW}$, where we denote by $\mathbf{GW}$ the unramified Grothendieck-Witt sheaf. On the other hand, if k is an infinite perfect field of characteristic unequal 2, then by [AH11, Lem. 4.4.4 & Thm. 4.4.5] the structure map $\mathbf{H}_0^{st\mathbb{A}^1}(X) \rightarrow \mathbf{H}_0^{st\mathbb{A}^1}(\operatorname{Spec} k)$ is a split epimorphism if and only if X has a zero cycle of degree one, where X is a smooth proper scheme over k . So if X is a smooth proper stable $\mathbb{A}^1$ -connected k -scheme, then the index of X over k is necessarily equal to 1. Of course, the existence of a rational point over k will imply the existence of a zero cycle of degree one over k . Now we state the following result:

Theorem 1.3. *Let k be an infinite perfect field of $\operatorname{char}(k) \neq 2$. Let X be a smooth projective stable $\mathbb{A}^1$ -connected k -scheme. If $U \subset X$ is an open subscheme such that $\operatorname{codim}(X \setminus U) \geq 2$, then U is also stable $\mathbb{A}^1$ -connected.*

2. A SIMPLE CHARACTERIZATION OF STABLE $\mathbb{A}^1$ -CONNECTEDNESS

We fix an infinite field k . Very similar to $\mathcal{SH}(k)$, whose homological t -structure is described by F. Morel in [Mor04], the t -structure of $D_{\mathbb{A}^1}(k)$ is given in [AH11, Thm. 3.3.3], which is the category of homotopy modules. Recall (cf. [AH11, Defn. 3.3.1]) that a homotopy module is a pair (F_*, ε_*) , where F_* is a $\mathbb{Z}$ -graded strictly $\mathbb{A}^1$ -invariant sheaf and for each $n \in \mathbb{Z}$ one has an isomorphism of abelian sheaves

$$\varepsilon_n : F_n \rightarrow (F_{n+1})_{-1}.$$

One denotes by $\mathcal{Ab}^{st\mathbb{A}^1}(k)$ the category of homotopy modules, whose morphisms are homogeneous natural transformations of $\mathbb{Z}$ -graded sheaves compatible with the isomorphisms above. One has

Proposition 2.1. [AH11, Prop. 3.4.1] *Let $F_* \in \mathcal{Ab}^{st\mathbb{A}^1}(k)$. One has canonical bijection contravariantly functorial in $\mathcal{X}$:*

$$H_{Nis}^0(\mathcal{X}, F_0) \xrightarrow{\cong} \operatorname{Hom}_{\mathcal{Ab}^{st\mathbb{A}^1}(k)}(\mathbf{H}_0^{st\mathbb{A}^1}(\mathcal{X})_*, F_*).$$

This implies immediately the following

Proposition 2.2. *Let $\mathcal{X}$ be a k -space. $\mathcal{X}$ is stable $\mathbb{A}^1$ -connected if and only if for any $F_* \in \mathcal{Ab}^{st\mathbb{A}^1}(k)$, one has a bijection*

$$H_{Nis}^0(\mathcal{X}, F_0) \cong \operatorname{Hom}_{\mathcal{Ab}^{st\mathbb{A}^1}}(\mathbf{GW}_*, F_*)$$

Example 2.3. If X/k is a smooth projective k -rational variety, then X is stable $\mathbb{A}^1$ -connected. Indeed, by [AH11, Prop. 3.4.2], the sheaf $\mathbf{H}_0^{st\mathbb{A}^1}(X)$ is a birational invariance in sense that if X and X' are birationally equivalent smooth proper k -varieties, then $\mathbf{H}_0^{st\mathbb{A}^1}(X) \cong \mathbf{H}_0^{st\mathbb{A}^1}(X')$. Thus, it is enough to see that $\mathbf{H}_0^{st\mathbb{A}^1}(\mathbb{P}_k^n) \cong \mathbf{H}_0^{st\mathbb{A}^1}(\mathbf{1}_k)$. This follows immediately from the distinguished triangle

$$C_*^{st\mathbb{A}^1}(\mathbb{P}^n) \rightarrow C_*^{st\mathbb{A}^1}(\mathbb{P}^{n+1}) \rightarrow \mathbb{Z}(n+1)[2n+2] \rightarrow C_*^{st\mathbb{A}^1}(\mathbb{P}^n)[1]$$

Remark 2.4. The same proof as in [A12, Thm. 3.9], one can even show that $\mathbf{H}_0^{st\mathbb{A}^1}(-)$ is a stably k -birational invariance of smooth proper k -varieties, i.e. if X is stably k -birationally equivalent to X' , then $\mathbf{H}_0^{st\mathbb{A}^1}(X) \cong \mathbf{H}_0^{st\mathbb{A}^1}(X')$. So as in [N12, Thm. 1.4], $\mathbf{H}_0^{st\mathbb{A}^1}(X)$ yields a

well-defined invariant on the Grothendieck ring $K_0(\text{Var}/k)/\mathbf{L}$, where $\mathbf{L}$ denotes the Lefschetz class $[\mathbb{A}_k^1]$.

3. PROOF OF 1.3

Throughout this section, we fix a base field k , which is an infinite perfect field of $\text{char}(k) \neq 2$. Let X be a smooth projective stable $\mathbb{A}^1$ -connected k -scheme. We write $Z = X \setminus U$ for the complement and $c = \text{codim}_X(Z) \geq 2$. Assume first of all Z is smooth, remark that we can use the stable $\mathbb{A}^1$ -connectivity theorem (cf. [Mor05]) of F. Morel to prove immediately the theorem, however we give here in the first part another proof including duality and calculation with $\mathbb{A}^1$ -cohomology. Consider then the Gysin triangle in $D_{\mathbb{A}^1}(k)$ (see [Deg08])

$$C_*^{st\mathbb{A}^1}(U) \rightarrow C_*^{st\mathbb{A}^1}(X) \rightarrow C_*^{st\mathbb{A}^1}(Z)(c)[2c] \rightarrow C_*^{st\mathbb{A}^1}(U)[1].$$

This yields an exact sequence of abelian groups

$$\begin{aligned} \cdots \rightarrow \text{Hom}_{D_{\mathbb{A}^1}(k)}(C_*^{st\mathbb{A}^1}(\text{Spec } L), C_*^{st\mathbb{A}^1}(Z)(c)[2c-1]) &\rightarrow \text{Hom}_{D_{\mathbb{A}^1}(k)}(C_*^{st\mathbb{A}^1}(\text{Spec } L), C_*^{st\mathbb{A}^1}(U)) \\ \rightarrow \text{Hom}_{D_{\mathbb{A}^1}(k)}(C_*^{st\mathbb{A}^1}(\text{Spec } L), C_*^{st\mathbb{A}^1}(X)) &\rightarrow \text{Hom}_{D_{\mathbb{A}^1}(k)}(C_*^{st\mathbb{A}^1}(\text{Spec } L), C_*^{st\mathbb{A}^1}(Z)(c)[2c]) \rightarrow \cdots \end{aligned}$$

where $L \in \mathcal{F}_k$ is a separable finitely generated field extension over k . By definition, we have $\text{Hom}_{D_{\mathbb{A}^1}(k)}(C_*^{st\mathbb{A}^1}(\text{Spec } L), C_*^{st\mathbb{A}^1}(X)) = \mathbf{H}_0^{st\mathbb{A}^1}(X)(L)$ and analogously for U . Now we prove that the other two terms in the exact sequence above are trivial. By Atiyah-Spanier-Whitehead duality in $D_{\mathbb{A}^1}(k)$ (see [AH11, §3.5]), we can write

$$\text{Hom}_{D_{\mathbb{A}^1}(k)}(C_*^{st\mathbb{A}^1}(\text{Spec } L), C_*^{st\mathbb{A}^1}(Z)(c)[2c-1]) \cong H_{st\mathbb{A}^1}^{2(c+n_Z+d_Z)-1}(Th(V_{Z_L}), \mathbb{Z}(c+n_Z+d_Z)),$$

where the vector bundle V_Z comes from [Voe03, Thm. 2.11] and n_Z denotes its rank. Here we write $d_Z = \dim(Z)$ and $H_{st\mathbb{A}^1}^*(-, \mathbb{Z}(*))$ is the $\mathbb{A}^1$ -stable cohomology. We compute first of all the unstable $\mathbb{A}^1$ -cohomology $H_{\mathbb{A}^1}^{2(c+n_Z+d_Z)-1}(Th(V_{Z_L}), \mathbb{Z}(c+n_Z+d_Z))$. By [AH11, Prop. 3.2.5], it is nothing but the hypercohomology $\mathbb{H}_{Nis}^*(-, \mathbb{Z}_{\mathbb{A}^1}(*))$. Consider the hypercohomology spectral sequence

$$E_2^{p,q} = H_{Nis}^p(-, \underline{H}^q(\mathbb{Z}_{\mathbb{A}^1}(*))) \Rightarrow \mathbb{H}_{Nis}^{p+q}(-, \mathbb{Z}_{\mathbb{A}^1}(*)).$$

As $\underline{H}^q(\mathbb{Z}_{\mathbb{A}^1}(*)) = 0$ for $q > *$ (a consequence of Morel's stable $\mathbb{A}^1$ -connectivity theorem) and $\mathbf{K}_*^{KW} = \underline{H}^*(\mathbb{Z}_{\mathbb{A}^1}(*))$ for $* > 0$ (a consequence of stable Hurewicz theorem), we have a canonical homomorphism

$$(3.1) \quad H_{\mathbb{A}^1}^{2(c+n_Z+d_Z)}(Th(V_{Z_L}), \mathbb{Z}(c+n_Z+d_Z)) \rightarrow H_{Nis}^{c+d_Z+n_Z-1}(Th(V_{Z_L}), \mathbf{K}_{c+n_Z+d_Z}^{MW}).$$

Since $c \geq 2$, which implies that $2(c+n_Z+d_Z) - 1 > cd_{Nis}(Th(V_{Z_L})) + c + n_Z + d_Z$ (remark that $cd_{Nis}(Th(V_{Z_L})) \leq n_Z + d_Z$, cf. [AH11, Lem. 3.1.7]), so by [AH11, Cor. 3.2.6] the above homomorphism is an isomorphism. But the group in the right hand side of the homomorphism 3.1 is trivial because of cohomological dimension reason. So we see that the unstable $\mathbb{A}^1$ -cohomology $H_{\mathbb{A}^1}^{2(c+n_Z+d_Z)}(Th(V_{Z_L}), \mathbb{Z}(c+n_Z+d_Z))$ is trivial. Let $j > 0$ be a natural number, we consider the groups

$$H_{\mathbb{A}^1}^{2(c+n_Z+d_Z)+j-1}(Th(V_{Z_L}) \wedge \mathbb{G}_m^{\wedge j}, \mathbb{Z}(c+n_Z+d_Z+j)).$$

These groups can be rewritten as $H_{\mathbb{A}^1}^{2(c+n_Z+d_Z)+j-1}(Th(V_{Z_L}), \mathbb{Z}(c+n_Z+d_Z+j))_{-(j)}$. So the same argument with the spectral sequence of hypercohomology as above yields

$$H_{\mathbb{A}^1}^{2(c+n_Z+d_Z)+j-1}(Th(V_{Z_L}), \mathbb{Z}(c+n_Z+d_Z+j)) \cong H_{Nis}^{c+n_Z+d_Z-1}(Th(V_{Z_L}), \mathbf{K}_{c+n_Z+d_Z+j}^{MW}),$$

which are trivial again by cohomological dimension reason ($c \geq 2$). So we may conclude now that the stable $\mathbb{A}^1$ -cohomology $H_{st\mathbb{A}^1}^{2(c+n_Z+d_Z)-1}(Th(V_{Z_L}), \mathbb{Z}(c+n_Z+d_Z))$ is trivial. Similarly, the group $\text{Hom}_{D_{\mathbb{A}^1}(k)}(C_*^{st\mathbb{A}^1}(\text{Spec } L), C_*^{st\mathbb{A}^1}(Z)(c)[2c])$ must also vanish. We see then $\mathbf{H}_0^{st\mathbb{A}^1}(U)(L) \cong \mathbf{H}_0^{st\mathbb{A}^1}(X)(L) \cong GW(L)$. In general, if Z is not smooth, since k is perfect, there is a finite stratification of closed subschemes of Z

$$\emptyset \subset Z_{d_X} \subset Z_{d_X-1} \subset \cdots \subset Z_{c+1} \subset Z_c = Z,$$

such that each $Z_i \setminus Z_{i+1}$ is smooth in $X \setminus Z_{i+1}$. So we may prove recursively and come back to the case when $Z = Z_{d_X}$, which follows from the previous case, since 0-dimensional schemes are smooth. More precisely, let us consider the Gysin triangle

$$C_*^{st\mathbb{A}^1}(X - Z_i) \rightarrow C_*^{st\mathbb{A}^1}(X - Z_{i+1}) \rightarrow C_*^{st\mathbb{A}^1}(Z_i - Z_{i+1})(c_i)[2c_i],$$

where all $c_i \geq 2$. As before, we apply the functor $\text{Hom}_{D_{\mathbb{A}^1}(k)}(C_*^{st\mathbb{A}^1}(\text{Spec } L), -)$ and consider then the group $\text{Hom}_{D_{\mathbb{A}^1}(k)}(C_*^{st\mathbb{A}^1}(\text{Spec } L), C_*^{st\mathbb{A}^1}(Z_i - Z_{i+1})(c_i)[2c_i - j])$ for $j = 0, 1$. These groups can be rewritten as

$$\begin{aligned} & \text{Hom}_{D_{\mathbb{A}^1}(k)}(C_*^{st\mathbb{A}^1}(\text{Spec } L)[j - c_i], C_*^{st\mathbb{A}^1}(Z_i - Z_{i+1} \wedge \mathbb{G}_m^{\wedge c_i})) = \\ & = \text{colim}_n \text{Hom}_{D_{\mathbb{A}^1}^{eff}(k)}(\tilde{C}_*^{\mathbb{A}^1}(\text{Spec } L_+ \wedge \mathbb{G}_m^n)[j - c_i], \tilde{C}_*^{\mathbb{A}^1}((Z_i - Z_{i+1})_+ \wedge \mathbb{G}_m^{\wedge n+c_i})) = \\ & = \text{colim}_n H_{j-c_i}^{\mathbb{A}^1}((Z_i - Z_{i+1}) \wedge \mathbb{G}_m^{\wedge c_i+n})_{-(n)}(\text{Spec } L). \end{aligned}$$

The last group must vanishing according to the stable $\mathbb{A}^1$ -connectivity theorem of F. Morel [Mor05], as $c_i \geq 2$ and $j = 0, 1$.

REFERENCES

- [A12] A. Asok, Birational invariants and $\mathbb{A}^1$ -connectedness, *Journal für die reine und angewandte Mathematik*, DOI: 10.1515/crelle-2012-0034, (2012).
- [AD09] A. Asok, B. Dorant, $\mathbb{A}^1$ -homotopy groups, excision, and solvable quotients, *Adv. in Math.*, Vol 221, Iss. 4, 1144-1190, (2009).
- [AH11] A. Asok, C. Haesemeyer, The 0-th stable $\mathbb{A}^1$ -homotopy sheaf and quadratic zero cycles, arXiv:1108.3854v1 [math.AG], Preprint (2011).
- [Deg08] F. Déglise, Around the Gysin triangle II, *Doc. Math.*, **13**, 613-675, (2008).
- [Mor04] F. Morel, An introduction to $\mathbb{A}^1$ -homotopy theory, *Contemporary developments in algebraic K-theory*, ICTP Lect. Notes XV, Abdus Salam, Int. Cent. Theoret. Phys., Trieste, 357-441, (2004).
- [Mor05] F. Morel, The stable $\mathbb{A}^1$ -connectivity theorems, *K-theory*, **35**, 1-68, (2005).
- [MV01] F. Morel, V. Voevodsky, $\mathbb{A}^1$ -homotopy theory of schemes, *IHÉS Publ. Math.*, (**90**), 45-143, (2001).
- [N12] LDT. Nguyen, Unramified cohomology, $\mathbb{A}^1$ -connectedness, and the Chevalley-Waring problem in Grothendieck ring, *C. R. Acad. Sci. Paris, Ser. I* **350**, 613-615, (2012).
- [Voe03] V. Voevodsky, Motivic cohomology with $\mathbb{Z}/2$ -coefficients, *Publ. Math. IHÉS* (2003).